

AD-A130 503

A LOCAL EXISTENCE AND UNIQUENESS THEOREM FOR A  
K-BKZ-FLUID(U) WISCONSIN UNIV-MADISON MATHEMATICS  
RESEARCH CENTER M RENARDY JUN 83 MRC-TSR-2530

1/1

UNCLASSIFIED

DAAG29-80-C-0041

F/G 12/1

NL


END

DATE

FORMED

8 83

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A130503

MRC Technical Summary Report # 2530

A LOCAL EXISTENCE AND  
UNIQUENESS THEOREM FOR A K-BKZ-FLUID

Michael Renardy

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

June 1983

(Received April 6, 1983)

DTIC FILE COPY

DTIC  
ELECTE  
JUL 21 1983  
S  
E  
D

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

National Science Foundation  
Washington, DC 20550

88 07 20 038

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

A LOCAL EXISTENCE AND UNIQUENESS THEOREM  
FOR A K-BKZ-FLUID

Michael Renardy

Technical Summary Report #2530

June 1983

ABSTRACT

We consider the three-dimensional motion of a viscoelastic liquid occupying all space. The constitutive law is assumed to be of the form suggested by Kaye [13] and Bernstein, Kearsley and Zapas [2]. An existence and uniqueness result for solutions of the initial value problem on sufficiently short time intervals is proved using Kato's theory of quasilinear hyperbolic equations.

AMS (MOS) Subjection Classifications: 35L15, 45K05, 47D05, 76A10

Key Words: Quasilinear Hyperbolic Equations, Viscoelastic Liquids, Semigroup theory

Work Unit Number 1 - Applied Analysis

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-7927062, Mod 2.

# SIGNIFICANCE AND EXPLANATION

The existence theory for models of viscoelastic fluids has so far not been very well developed, in particular in three dimensional situations. <sup>the author</sup> Here, ~~we~~ prove an existence theorem for a particular class of models, suggested by Kaye and Bernstein, Kearsley and Zapas. This theory is based on a postulated analogy with hyperelasticity. It is assumed that the fluid occupies all of space. Abstract methods developed originally for quasilinear hyperbolic systems can be used to prove the well-posedness of the initial value problem.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

# A LOCAL EXISTENCE AND UNIQUENESS THEOREM FOR A K-BKZ-FLUID

Michael Renardy

## 1. Introduction

The purpose of this paper is to show the well-posedness of the initial history value problem for a class of viscoelastic materials with memory. The constitutive law is assumed to be of the form proposed by Kaye [13] and Bernstein, Kearsley and Zapas [2]. Their theory is motivated by a formal analogy with classical hyperelasticity [21]. The equilibrium configuration is replaced by a configuration previously occupied by the material. It is assumed that the stress contributions from all the previous configurations superpose in an additive fashion, leading mathematically to a convolution integral.

We consider a material occupying all of  $\mathbb{R}^3$ , which is assumed incompressible. By  $\underline{\zeta} = (\zeta^1, \zeta^2, \zeta^3)$  we denote body coordinates, while  $\underline{y}(\underline{\zeta}, t) = (y^1, y^2, y^3)$  denotes the position of material points in space. The equation of motion is given by

$$\begin{aligned} \rho y^i &= \frac{\partial}{\partial \zeta^s} \left( \frac{\partial y^i}{\partial \zeta^r} \pi^{rs} \right) + g^i(\underline{\zeta}, t) \\ (1.1) \quad \det \left( \frac{\partial y^i}{\partial \zeta^j} \right) &= 1. \end{aligned}$$

Here  $\underline{\pi}$  denotes the convected stress tensor, and the  $g^i$  are the components of the body force (we could allow  $g^i$  to depend on  $\underline{y}$ , but omit this for simplicity). The Einstein convention is used throughout this paper.

We also need the following notations:

$$\underline{\gamma} = (\gamma_{ij}) = \left( \frac{\partial y^i}{\partial \zeta^j} \right), \quad \gamma^{ij} \gamma_{jk} = \delta^i_k$$

$$I_1(\underline{\zeta}, t, \tau) = \text{tr}(\underline{\gamma}(\underline{\zeta}, t) \underline{\gamma}^{-1}(\underline{\zeta}, \tau))$$

$$I_2(\underline{x}, t, \tau) = \text{tr}(\underline{\gamma}^{-1}(\underline{x}, t) \underline{\gamma}(\underline{x}, \tau)) .$$

The K-BKZ constitutive model is described in terms of a function  $W$ , which is the analogue of the strain energy in classical hyperelasticity.  $W$  is a real-valued function of the two arguments  $I_1$  and  $I_2$ . The constitutive law of K-BKZ is

$$\begin{aligned} \tau^{rs} = & -p\gamma^{rs} + \int_{-\infty}^t a(t-\tau) \left[ \frac{\partial W}{\partial I_1} \cdot \gamma^{rs}(\tau) \right. \\ (1.2) \quad & \left. - \frac{\partial W}{\partial I_2} \gamma^{rp}(\tau) \gamma_{pq}(\tau) \gamma^{qs}(\tau) \right] d\tau . \end{aligned}$$

Here  $p$  denotes the undetermined pressure. The kernel  $a$  is positive and monotone decreasing. We also assume that  $W$  and  $a$  are smooth functions and that  $a$  decays sufficiently fast (e.g., exponentially) at infinity. In particular, we do not admit that  $a$  is singular at 0. Such singularities, however, are physically possible and are in fact predicted by some molecular theories, e.g. [5]. Thus the requirement that  $a$  is smooth must be regarded as a restriction on the material.

The K-BKZ theory includes various models suggested in the literature - some of them based on molecular theories - as special cases [3], [5], [7], [15-17], [22], [23]. A crucial hypothesis needed for well-posedness of the initial value problem is that the equation of motion is "hyperbolic", or, equivalently, that the corresponding quasi-static equation is "elliptic". This requirement leads to an inequality involving first and second derivatives of  $W$ . In chapter 2, we give a sufficient condition for ellipticity. We show that ellipticity holds if  $W$  is monotone in each argument and is a convex function of  $\sqrt{I_1}$  and  $\sqrt{I_2}$ . These conditions imply in particular that, in simple shear flow,  $N_1$  is positive,  $N_2$  is negative and  $|N_1| > |N_2|$ . These consequences are consistent with experimental results on polymer melts and solutions.

Our well-posedness theorem is based on Kato's theory of quasilinear hyperbolic systems [8], [10-12]. The case of classical hyperelasticity is considered in [8], and, to some extent, our arguments will be analogous. However, the history dependence and

incompressibility conditions introduce complicating features. As a result, Kato's assumptions cannot be verified immediately, using the equations as they stand, but only after going through some transformations and substitutions.

We shall not consider (1.1) itself, but its second time-derivative. Equation (1.1) and its first time derivative are then used to express  $\dot{\gamma}$  and  $\ddot{\gamma}$  in terms of  $\ddot{\gamma}$  and  $\dddot{\gamma}$ . This elimination is discussed in chapter 3. Because of ellipticity, there is a gain of regularity associated with it. Roughly speaking, this serves to improve the smoothness of the "coefficients" in a quasilinear formulation of the problem.

In chapter 4, we finally pose the evolution problem in a form accessible to Kato's method. The delay equation is reformulated as an abstract evolution problem on a history space. In contrast with some previous approaches to similar problems, however, this history space only involves the history back to the initial time  $t = 0$  rather than all the way back to  $t = -\infty$ . In the previous literature, existence theorems for "hyperbolic" models of viscoelastic fluids mostly concern linearized [9], [20] or nonlinear, but one-dimensional problems [4], [18]. The present problem has been studied by Kim [14] in the special case  $W = I_2$ . Kim's method is based on artificial viscosity and differs substantially from the approach used here.



## 2. A sufficient condition for ellipticity

For abbreviation, let  $\tilde{F}$  be defined by  $\tilde{F}_j^i = \frac{\partial y^i}{\partial z^j}$ , and let  $F(t, \tau) = \tilde{F}(t) \tilde{F}^{-1}(\tau) = \frac{\partial y(t)}{\partial y(\tau)}$ . Then  $I_1 = \text{tr}(F^T F)$ , and  $I_2 = \text{tr}(F^{-1} F^{-T})$ , so that  $W$  is a function of  $F$ . The right hand side of (1.1) can be rewritten as follows

$$(2.1) \quad \frac{\partial}{\partial z^s} \left( \frac{\partial y^i}{\partial z^r} \pi^{rs} \right) = - \frac{\partial p}{\partial z^s} \frac{\partial z^s}{\partial y^i} + \frac{1}{2} \int_{-\infty}^t a(t-\tau) \cdot$$

$$\frac{\partial^2 W}{\partial F_p^i \partial F_r^j} \left[ \frac{\partial^2 y^j}{\partial z^q \partial z^s} \frac{\partial z^q}{\partial y^r(\tau)} \frac{\partial z^s}{\partial y^p(\tau)} + \frac{\partial y^j}{\partial z^q} \right.$$

$$\left. \frac{\partial}{\partial z^s} \left( \frac{\partial z^q}{\partial y^r(\tau)} \right) \frac{\partial z^s}{\partial y^p(\tau)} \right] d\tau.$$

Let  $\bar{F}_i^p = \frac{\partial y^p(\tau)}{\partial y^i(t)}$  denote the entries of  $F^{-1}$ . The right hand side of (2.1) remains

unchanged if  $\frac{\partial^2 W}{\partial F_p^i \partial F_r^j}$  is replaced by  $\frac{\partial^2 W}{\partial F_p^i \partial F_r^j} + K \bar{F}_i^p \bar{F}_j^r$ . The additional term vanishes as a result of the incompressibility condition. In the subsequent chapters, we shall assume that the following strong ellipticity condition is satisfied:

(E). If  $K > 0$  is chosen large enough, then

$$\left( \frac{\partial^2 W}{\partial F_p^i \partial F_r^j} + K \bar{F}_i^p \bar{F}_j^r \right) \lambda^i \lambda^j \mu_p \mu_r > c |\lambda|^2 |\mu|^2, \quad c > 0.$$

In terms of the original arguments of  $W$ ,  $I_1$  and  $I_2$ , this condition has a rather indirect form. However, we have the following sufficient criterion, which is easily expressed in terms of  $I_1$  and  $I_2$ .

### Lemma 2.1:

If  $W$  is monotone in each argument, strictly monotone in  $I_1$  or  $I_2$ , and a convex function of  $\sqrt{I_1}$  and  $\sqrt{I_2}$ , then (E) holds.

Proof:

We have

$$\begin{aligned}
\frac{\partial^2 W}{\partial F_p^i \partial F_r^j} &= 2 \frac{\partial W}{\partial I_1} \delta_{ij} \delta^{pr} \\
&+ 2 \frac{\partial W}{\partial I_2} (\bar{F}_j^p \bar{F}_i^r \bar{F}_\sigma^p \bar{F}_\sigma^r \\
&\quad + \bar{F}_i^p \bar{F}_j^r \bar{F}_\sigma^p \bar{F}_\sigma^r \\
&\quad + \bar{F}_i^p \bar{F}_\sigma^p \bar{F}_j^r \bar{F}_\sigma^r) \\
&+ 4 \frac{\partial^2 W}{\partial I_1^2} F_p^i F_r^j \\
&- 4 \frac{\partial^2 W}{\partial I_1 \partial I_2} (\bar{F}_p^i \bar{F}_j^p \bar{F}_\sigma^r \bar{F}_\sigma^p + \bar{F}_r^j \bar{F}_i^p \bar{F}_\sigma^p \bar{F}_\sigma^p) \\
&+ 4 \frac{\partial^2 W}{\partial I_2^2} (\bar{F}_i^p \bar{F}_\sigma^p \bar{F}_\sigma^p \bar{F}_j^r \bar{F}_\sigma^r \bar{F}_\sigma^r) .
\end{aligned}$$

Moreover,  $\frac{\partial W}{\partial \sqrt{I_1}} = 2\sqrt{I_1} \frac{\partial W}{\partial I_1}$

$$\frac{\partial^2 W}{(\partial \sqrt{I_1})^2} = 2 \frac{\partial W}{\partial I_1} + 4 I_1 \frac{\partial^2 W}{\partial I_1^2} .$$

Using this, we obtain

$$\begin{aligned}
\frac{\partial^2 W}{\partial F_p^i \partial F_r^j} &= \frac{\partial W}{\partial \sqrt{I_1}} \left[ \frac{1}{\sqrt{I_1}} \delta_{ij} \delta^{pr} - \frac{1}{I_1^{3/2}} F_p^i F_r^j \right] \\
&+ \frac{\partial W}{\partial \sqrt{I_2}} \cdot \left[ \frac{1}{\sqrt{I_2}} (\bar{F}_j^p \bar{F}_i^r \bar{F}_\sigma^p \bar{F}_\sigma^r + \bar{F}_i^p \bar{F}_j^r \bar{F}_\sigma^p \bar{F}_\sigma^r \right. \\
&\quad \left. + \bar{F}_i^p \bar{F}_\sigma^p \bar{F}_j^r \bar{F}_\sigma^r) \right. \\
&\quad \left. - \frac{1}{I_2^{3/2}} (\bar{F}_i^p \bar{F}_\sigma^p \bar{F}_\sigma^p \bar{F}_j^r \bar{F}_\sigma^r \bar{F}_\sigma^r) \right] \\
&+ \frac{1}{I_1} \frac{\partial^2 W}{(\partial \sqrt{I_1})^2} F_p^i F_r^j
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{\sqrt{I_1 I_2}} \frac{\partial^2 W}{\partial \sqrt{I_1} \partial \sqrt{I_2}} (\bar{F}_p^i \bar{F}_j^p \bar{F}_\sigma^r \bar{F}_\sigma^p + \bar{F}_r^j \bar{F}_i^p \bar{F}_\sigma^p \bar{F}_\sigma^p) \\
& + \frac{1}{I_2} \frac{\partial^2 W}{(\partial \sqrt{I_2})^2} (\bar{F}_i^p \bar{F}_\sigma^p \bar{F}_\sigma^p \bar{F}_j^u \bar{F}_v^r \bar{F}_v^u) .
\end{aligned}$$

The first term leads to  $\frac{\partial W}{\partial \sqrt{I_1}} \frac{1}{\sqrt{I_1}} [|\lambda|^2 |\mu|^2 - \frac{1}{I_1} (\underline{\mu} \cdot \underline{\underline{g}} \underline{\lambda})^2] > 0$ . The last three terms yield

a positive contribution, if  $W$  is convex. To estimate the second term, let us put

$\bar{F}_i^p \lambda^i = \tilde{\lambda}^p$ ,  $\bar{F}_\sigma^p \bar{F}_\sigma^p = \alpha^{pp}$ . We find

$$\begin{aligned}
& \frac{\partial W}{\partial \sqrt{I_2}} \cdot \frac{1}{\sqrt{I_2}} (\tilde{\lambda}^p \tilde{\lambda}^r \alpha^{pp} \mu_p^u \mu_r + \tilde{\lambda}^p \tilde{\lambda}^p \alpha^{pr} \mu_p^u \mu_r \\
& + \tilde{\lambda}^p \tilde{\lambda}^p \alpha^{pr} \mu_p^u \mu_r - \frac{1}{I_2} \tilde{\lambda}^p \tilde{\lambda}^u \alpha^{pp} \alpha^{ru} \mu_p^u \mu_r) \\
& = \frac{\partial W}{\partial \sqrt{I_2}} \cdot \frac{1}{\sqrt{I_2}} [2(\tilde{\lambda} \cdot \underline{\mu})(\tilde{\lambda} \cdot \underline{\underline{g}} \cdot \underline{\mu}) + |\tilde{\lambda}|^2 (\underline{\mu} \cdot \underline{\underline{g}} \cdot \underline{\mu}) - \frac{1}{I_2} (\tilde{\lambda} \cdot \underline{\underline{g}} \cdot \underline{\mu})^2] .
\end{aligned}$$

The term  $K \bar{F}_i^p \bar{F}_j^r$  gives a contribution  $K \cdot (\tilde{\lambda} \cdot \underline{\mu})^2$ . Since  $I_2 = \text{tr } \alpha$ , we have

$|\tilde{\lambda}|^2 (\underline{\mu} \cdot \underline{\underline{g}} \cdot \underline{\mu}) - \frac{1}{I_2} (\tilde{\lambda} \cdot \underline{\underline{g}} \cdot \underline{\mu})^2 > 0$ , and the whole term can be made positive by choosing  $K$  large enough. This proves the lemma.

### 3. Solution of a nonlinear elliptic Volterra equation

Throughout the remainder of the paper, it is assumed that (E) holds with a suitable constant  $K \in \mathbb{R}$ . For abbreviation, we write  $\Lambda_{ij}^{pr} = \frac{\partial^2 W}{\partial F_i^p \partial F_j^r} + K F_i^p F_j^r$ . Thus  $\Lambda_{ij}^{pr}$  depends on the gradients of  $y(t)$  and  $y(\tau)$ , and it is symmetric in  $(i,p)$  and  $(j,r)$ . Equation (1.1) is rewritten in the following equivalent form

$$\begin{aligned} h^i &:= \rho \ddot{y}^i - \lambda(y^i - \zeta^i) - g^i = \\ &= - \frac{\partial p}{\partial \zeta^s} \frac{\partial \zeta^s}{\partial y^i} + \frac{1}{2} \int_{-\infty}^t a(t-\tau) \Lambda_{ij}^{pr} \cdot \frac{\partial}{\partial \zeta^s} \\ &\quad \left[ \frac{\partial y^j}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial y^r(\tau)} \frac{\partial \zeta^s}{\partial y^p(\tau)} \right] d\tau - \lambda(y^i - \zeta^i) \\ 0 &= \det \left( \frac{\partial y^i}{\partial \zeta^j} \right) - 1. \end{aligned} \quad (3.1)$$

It is convenient to split the history occurring in the integral into a known and an unknown part. Let the initial time be  $t = 0$ . Then  $y(\tau)$ ,  $\tau < 0$  is considered known, and we shall denote it by  $y_0(\tau)$ . On the interval  $[0, t]$ , we perform a rescaling:  $\hat{y}(\sigma) = y(\sigma t)$ , so that  $\sigma$  ranges from 0 to 1. Equation (3.1) can be rewritten as an equation for  $\hat{y}$  as follows:

$$\begin{aligned} \hat{h}^i(\sigma) &= - \frac{\partial p(\sigma)}{\partial \zeta^s} \frac{\partial \zeta^s}{\partial \hat{y}^i(\sigma)} + \frac{1}{2} \int_{-\infty}^0 a(\sigma t - \tau) \cdot \\ &\quad \Lambda_{ij}^{pr}(\hat{y}(\sigma), y_0(\tau)) \frac{\partial}{\partial \zeta^s} \left[ \frac{\partial \hat{y}^j(\sigma)}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial y_0^r(\tau)} \frac{\partial \zeta^s}{\partial y_0^p(\tau)} \right] d\tau \\ &\quad + \frac{1}{2} t \int_0^\sigma a(\sigma t - \sigma' t) \Lambda_{ij}^{pr}(\hat{y}(\sigma), \hat{y}(\sigma')) \cdot \frac{\partial}{\partial \zeta^s} \\ &\quad \left[ \frac{\partial \hat{y}^j(\sigma)}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial \hat{y}^r(\sigma')} \frac{\partial \zeta^s}{\partial \hat{y}^p(\sigma')} \right] d\sigma' - \lambda(\hat{y}^i(\sigma) - \zeta^i) \\ 0 &= \det \left( \frac{\partial \hat{y}^i(\sigma)}{\partial \zeta^j} \right) - 1. \end{aligned} \quad (3.2)$$

The term involving the unknown part of the history (from 0 to  $t$ ) is now multiplied by a factor of  $t$ . Hence it can be treated as a perturbation for  $t$  small enough. If we set  $t = 0$ , then the right hand side of (3.2) acts pointwise with respect to  $\sigma$  and, for each  $\sigma \in [0,1]$ , we have a nonlinear elliptic system for  $\hat{y}(\sigma)$  and  $\hat{p}(\sigma)$ .

We assume that we have a solution  $\hat{y}(\sigma) \equiv y(0)$ ,  $\hat{p}(\sigma) \equiv p(0)$  for  $t = 0$  given by the initial conditions of the problem. If the linearization of the right hand side of (3.2) at this given solution is invertible, then the implicit function theorem can be used to show the existence of solutions to (3.2) for small  $t$ .

More precisely, let  $n$  be any integer  $> 1$ . We assume that the initial history is such that  $y_0(\tau) - \underline{z}$  is continuous and bounded from  $(-\infty, 0]$  into  $H^{n+2}(\mathbb{R}^3)$  and that it satisfies the incompressibility condition. Let the initial values  $\underline{h}(0) \in H^n$ ,  $y(0) \in \underline{z} + H^{n+2}$  and  $\nabla p(0) \in H^n$  satisfy (3.1) for  $t = 0$ . Then the constant functions  $\hat{h}(\sigma) \equiv \underline{h}(0)$  etc. satisfy (3.2) for  $t = 0$ . Condition (E) guarantees that the linearization of the right hand side of (3.2) (for  $t = 0$ ) is an elliptic system in the sense of Agmon, Douglis and Nirenberg [1], [6], moreover, it is invertible if  $\lambda > 0$  is chosen large enough. The existence of a weak solution is established by variational methods using Gårding's inequality. Regularity can then be obtained using the standard trick of estimating difference quotients. According to the implicit function theorem, equation (3.2) can be solved uniquely for  $\hat{y} \in C([0,1]; \underline{z} + H^{n+2})$ ,  $\hat{p} \in C([0,1]; H^n)$  as functions of  $\hat{h} \in C([0,1]; H^n)$ , provided  $t$  is small enough. For technical reasons, we shall have to use a different topology for the  $\sigma$ -dependence in chapter 4, but a similar argument will apply also with that topology.

By applying the same procedure to the time differentiated version of (1.1), we can resolve for  $\dot{\hat{y}}$  and  $\dot{\hat{p}}$  as functions of  $\dot{\hat{h}}$ ,  $\hat{y}$  and  $\hat{p}$ .

Remark: The reader should be careful to distinguish between  $\dot{\hat{y}}$  and  $\dot{\hat{y}}$ . We have  $\dot{\hat{y}}(\sigma) = \dot{\hat{y}}(\sigma t)$ , but  $\dot{\hat{y}}(\sigma) = \frac{d}{dt} \hat{y}(\sigma t) = \sigma \dot{\hat{y}}(\sigma t)$ .

#### 4. The hyperbolic Cauchy problem

We use the abbreviations:  $\underline{u} = \dot{\underline{y}}$ ,  $\underline{v} = \ddot{\underline{y}} - \frac{\lambda}{\rho} (\underline{y} - \underline{\zeta})$ ,  $\underline{w} = \ddot{\underline{y}} - \frac{\lambda}{\rho} \dot{\underline{y}}$ ,  $\underline{q} = \dot{\underline{p}}$ ,  $\phi = \ddot{\underline{p}}$ , where  $\lambda$  is as in the previous chapter. The second time derivative of equation (1.1) can then be written as follows:

$$\begin{aligned} \dot{\underline{v}} &= \underline{w} \\ \rho \dot{w}^i &= - \frac{\partial \phi}{\partial \zeta^s} \frac{\partial \zeta^s}{\partial y^i} + \frac{1}{2} \int_{-\infty}^t a(t-\tau) \lambda_{ij}^{pr} \frac{\partial}{\partial \zeta^s} \\ (4.1) \quad & \left( \frac{\partial v^j}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial y^r(\tau)} \frac{\partial \zeta^s}{\partial y^p(\tau)} \right) d\tau + \ddot{g}^i \\ & + \phi^i \{ \underline{y}, \underline{u}, \underline{v}, \underline{p}, \underline{q} \} \\ & \frac{\partial w^i}{\partial \zeta^j} \frac{\partial \zeta^j}{\partial y^i} = \Psi(\underline{y}, \underline{u}, \underline{v}) . \end{aligned}$$

Here  $\phi$  is an expression involving spatial derivatives of  $\underline{y}$  and  $\underline{u}$  up to the second order, and spatial derivatives of  $\underline{v}$ ,  $\underline{p}$  and  $\underline{q}$  up to the first order.  $\Psi$  depends on first spatial derivatives of  $\underline{y}$ ,  $\underline{u}$  and  $\underline{v}$ . Thus the  $\phi$ - and  $\Psi$ -terms represent perturbations of lower differential order than the other terms in the equation. Of course, we think of  $\underline{y}$ ,  $\underline{u}$ ,  $\underline{p}$  and  $\underline{q}$  as being expressed in terms of  $\underline{v}$  and  $\underline{w}$  as described in section 3.

In order to represent (4.1) as an evolution problem, we have to eliminate the variable  $\phi$ . As in Navier-Stokes theory, this is done by using the Hodge projection. The difference is that, since we work in Lagrangian coordinates, the Hodge projection is a function of the displacement  $\underline{y}$ . Let  $\underline{y} \in \underline{\zeta} + H^{n+1}$  ( $n \geq 2$ ) be such that its gradient satisfies the incompressibility condition. We define an orthogonal projection  $P(\underline{y})$  in  $L^2(\mathbb{R}^3)$  to be such that its range consists of all vectorfields  $\underline{u} \in L^2$  which satisfy  $\frac{\partial u^i}{\partial \zeta^j} \frac{\partial \zeta^j}{\partial y^i} = 0$ , while its nullspace contains all vectorfields of the form  $\frac{\partial \zeta^s}{\partial y^i} \frac{\partial q}{\partial \zeta^s}$ , where  $\nabla q \in L^2$ . In other words,  $P(\underline{y})$  is the image of the Hodge projection under the coordinate transformation  $\underline{y} \mapsto \underline{\zeta}$ . The projection  $P(\underline{y})$  is continuous from  $H^j$  into itself for  $j \leq n$ , and it

can be shown [19] that  $P(y)$  - as an operator in  $H^j$  - depends smoothly on  $y \in \mathbb{R}^n + H^{n+1}$ .

Obviously,  $\phi$  is eliminated from (4.1) by applying the projection  $P(y)$ . Moreover, the incompressibility condition requires that

$$\frac{\partial v^i}{\partial \zeta^s} \frac{\partial \zeta^s}{\partial y^i} = \frac{\partial u^i}{\partial \zeta^s} \frac{\partial \zeta^s}{\partial y^k} \frac{\partial u^k}{\partial \zeta^l} \frac{\partial \zeta^l}{\partial y^i}.$$

This can be used to express  $(1 - P(y))v$  in terms of  $y$  and  $u$ . More precisely,  $(1 - P(y))v \in H^{n+2}$  is a smooth function of  $u \in H^{n+2}$ ,  $y \in H^{n+3}$ . Similarly, by differentiating the last equation of (4.1) with respect to time, we can express  $(1 - P(y))\dot{w} \in H^n$  as a smooth function of  $u, v, \dot{w} \in H^n$ ,  $y \in H^{n+1}$ . Thus we can insert or omit terms involving  $(1 - P(y))v$ ,  $(1 - P(y))\dot{w}$  and compensate for this by other terms involving only lower order derivatives.

We use this to write (4.1) in the following form

$$\begin{aligned} \dot{v} &= \underline{w} \\ \rho \dot{w} &= \frac{1}{2} P(y) \underline{A} P(y) v + (1 - P(y)) \Delta \\ &\quad (1 - P(y)) v + \hat{\Phi}(y, u, v, w, p, q) \\ &\quad - P(y) \ddot{q}. \end{aligned} \quad (4.2)$$

Here  $\underline{A}$  stands for

$$(\underline{A} v)^i = \frac{1}{2} \int_{-\infty}^t a(t-\tau) A_{ij}^{pr} \frac{\partial}{\partial \zeta^s} \left\{ \frac{\partial v^j}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial y^r(\tau)} \frac{\partial \zeta^s}{\partial y^p(\tau)} \right\} d\tau$$

and  $\Delta$  denotes the Laplace operator. The term  $(1 - P(y)) \Delta(1 - P(y))v$  will be convenient later. Of course, another term compensating for it is contained in  $\hat{\Phi}$ .

We rewrite (4.2) as an evolution problem on a history space by introducing the "hat variables" as in chapter 3:  $\hat{v}(\sigma) = v(\sigma t)$ ,  $\hat{w}(\sigma) = w(\sigma t)$ . The operator  $\underline{A}$  is split into two parts:

$$\begin{aligned} (\hat{A}_1(t, \hat{y}, \sigma) \hat{v}(\sigma))^i &= \frac{1}{2} t \int_{-\infty}^0 a(\sigma t - \tau) \hat{A}_{ij}^{pr}(\hat{y}(\sigma), \hat{y}_0(\tau)) \\ &\quad \frac{\partial}{\partial \zeta^s} \left\{ \frac{\partial \hat{v}^j(\sigma)}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial y_0^r(\tau)} \frac{\partial \zeta^s}{\partial y_0^p(\tau)} \right\} d\tau \end{aligned}$$

$$(\hat{A}_2(t, \hat{y}, \sigma) \hat{y}(\sigma))^i = \frac{1}{2} t \int_0^\sigma a(\sigma t - \sigma' t) \hat{A}_{ij}^{pr}(\hat{y}(\sigma), \hat{y}(\sigma'))$$

$$\frac{\partial}{\partial \zeta^s} \left[ \frac{\partial \hat{v}^j(\sigma)}{\partial \zeta^q} \frac{\partial \zeta^q}{\partial \hat{y}^r(\sigma')} \frac{\partial \zeta^s}{\partial \hat{y}^p(\sigma')} \right] d\sigma'.$$

Equation (4.2) can now be written in the following form:

$$(4.3) \quad \begin{aligned} \hat{v}(\sigma) &= \hat{\sigma} \hat{w}(\sigma) \\ \hat{w}(\sigma) &= \sigma \cdot \frac{1}{\rho} \frac{\{P(\hat{y}(\sigma))(\hat{A}_1 + \hat{A}_2)P(\hat{y}(\sigma))\hat{v}(\sigma) + (1 - P(\hat{y}(\sigma))\Delta(1 - P(\hat{y}(\sigma))\hat{v}(\sigma) + \hat{\phi}(\hat{v}, \hat{w})(\sigma) + P(\hat{y}(\sigma))\hat{q}(\sigma))\}}{\end{aligned}$$

It will turn out that (4.3) can be put into the context of an abstract result due to Hughes, Kato and Marsden [8]. In the following, we quote their assumptions and their result.

We are concerned with an evolution problem written in the form

$$(4.4) \quad \dot{u} = -A(t, u)u + f(t, u), \quad 0 < t < T, \quad u(0) = \phi,$$

where  $u$  lies in a Banach space. Since the operator  $A$  is in applications a differential operator, it does not take any Banach space into itself, and we have to consider more than one topology. We shall therefore be dealing with three Banach spaces:  $Y \subset X \subset Z$ , all reflexive and separable with continuous and dense inclusions. (Hughes, Kato and Marsden introduce a fourth space  $Z'$ , which for our application can be taken equal to  $X$ .)

On the space  $Z$ , a variable norm is considered. Let  $N(Z)$  be the set of all norms in  $Z$  equivalent to the given one  $\|\cdot\|_Z$ . A metric on  $N(Z)$  is given by

$$d(\|\cdot\|_\mu, \|\cdot\|_\nu) = \ln \max \left\{ \sup_{0 \neq z \in Z} \|z\|_\mu / \|z\|_\nu, \sup_{0 \neq z \in Z} \|z\|_\nu / \|z\|_\mu \right\}.$$

We now state the assumptions of Hughes, Kato and Marsden. All assumptions are supposed to hold for  $t, t', \dots \in [0, T]$  for some  $T > 0$  and  $w, w', \dots \in W$ , where  $W$  is an open set in  $Y$ .  $\beta, \lambda_N, \mu_N, \dots$  are arbitrary constants. The assumptions are:

(N) For  $(t, w) \in [0, T] \times W$ , there is  $N(t, w) \in N(Z)$  such that

$$d(N(t, w), \|\cdot\|_Z) < \lambda_N$$

$$d(N(t', w'), N(t, w)) < \mu_N(|t - t'| + \|w - w'\|_X).$$



(S) There is an isomorphism  $S(t,w) : Y \rightarrow Z$  such that

$$\|S(t,w)\|_{Y,Z} < \lambda_s, \|S^{-1}(t,w)\|_{Z,Y} < \lambda_s^{-1}$$

$$\|S(t',w') - S(t,w)\|_{Y,Z} < \mu_s(|t'-t| + \|w'-w\|_X).$$

(A1)  $A(t,w) \in G(Z_{N(t,w)}, 1, \beta)$ . This means that  $-A(t,w)$  is a  $C_0$ -generator in  $Z$

(equipped with the norm  $N(t,w)$ ) such that

$$\|e^{-\tau A(t,w)}\|_{Z_{N(t,w)}} < e^{\beta\tau} \|z\|_{N(t,w)}.$$

(A2)  $S(t,w)A(t,w)S^{-1}(t,w) = A(t,w) + B(t,w)$ , where

$$B(t,w) \in B(Z), \|B(t,w)\|_{Z,Z} < \lambda_B.$$

(A3)  $A(t,w) \in B(Y,X)$  with  $\|A(t,w)\|_{Y,X} < \lambda_A$

$$\text{and } \|A(t,w') - A(t,w)\|_{Y,X} < \mu_A \|w'-w\|_X.$$

The mapping  $t \mapsto A(t,w) \in B(Y,Z)$  is continuous in norm.

(A4) There is an element  $y_0 \in W$  such that

$$A(t,w)y_0 \in Y, \|A(t,w)y_0\| < \lambda_0.$$

(f1)  $f(t,w) \in Y$ ,  $\|f(t,w)\|_Y < \lambda_f$ ,  $\|f(t,w') - f(t,w)\|_X$

$$< \mu_f \|w'-w\|_X \text{ and the mapping } t \mapsto f(t,w) \in Z \text{ is continuous.}$$

Under these assumptions, the following theorem is proved.

**Theorem 4.1:**

Assume (N), (S), (A1) - (A4) and (f1) hold. Then there is  $\rho' > 0$  and  $0 < T' < T$  such that for  $\phi \in Y$  with  $\|\phi - y_0\|_Y < \rho'$ , equation (4.4) has a unique solution  $u$  on  $[0, T']$  such that  $u \in C([0, T']; W) \cap C^1([0, T']; X)$ . Here  $\rho'$  depends on  $\lambda_s, \lambda_s^{-1}, \lambda_N$  and  $R := \text{dist}(y_0, Y \setminus W)$ ,  $T'$  depends on all the constants occurring in the assumptions. When  $\phi$  varies in  $Y$  subject to the restriction  $\|\phi - y_0\| < \rho'$ , the mapping  $\phi \mapsto u(t)$  is Lipschitz continuous in the  $X$ -norm, uniformly for  $t \in [0, T']$ .

We now show how this applied to (4.3). For this, we have to identify the space  $Y$ ,  $X$  and  $Z$  and the functions  $A$ ,  $f$ ,  $S$  and  $N$ .

The definition of the spaces requires that we specify a topology both for the spatial dependence and for the  $\sigma$ -dependence. Both have to be reflexive. We denote by  $W^{0,n}$  the space of all functions  $\hat{y}(\sigma) : \sigma \in [0, 1] \rightarrow H^n(\mathbb{R}^3, \mathbb{R}^3)$ , which are square integrable over  $[0, 1]$  (in the Bochner sense).  $W^{1,n}$  denotes the space of all functions

$[0,1] \times H^n(\mathbb{R}^3, \mathbb{R}^3)$ , which have a square integrable derivative (with respect to  $\sigma$ ). Moreover, we let  $V^n = (W^{1,n+1} \cap W^{0,n+4}) \times (W^{1,n} \cap W^{0,n+3})$ . Our choice of spaces is  $Z = V^1$ ,  $X = V^n$ ,  $Y = V^{n+1}$ , where  $n > 4$ . The operator  $-A$  is identified with the underlined terms on the right hand side of (4.3), while  $f$  consists of the remaining terms. We take  $S$  to be equal to  $(\frac{\lambda}{\sigma} + \lambda)^n$ , where  $\lambda$  is such that this operator is invertible. (As in chapter 3, it follows from the theory of elliptic equations that such a  $\lambda$  exists.)

This last choice trivially satisfies (A2) with  $B = 0$ . Moreover, our choice of topologies is such that all boundedness and Lipschitz conditions are satisfied, (under appropriate smoothness assumptions on the kernel  $a$ , the strain energy  $W$  and the data of the problem) i.e. the conditions (S), (A3) and (f1). As in most applications, there is a dense set of  $y_0$  satisfying (A4) (any  $y_0 \in V^{n+2}$  does).

It thus remains to identify a norm  $N$  such that (A1) holds. For this, we write

$$-A = \sigma \begin{pmatrix} 0 & 1 \\ \lambda & 0 \end{pmatrix}.$$

The operator  $\hat{A}$  has a similar form as in [8], and the same argument as there shows that (A1) holds in the space  $W^{0,1} \times W^{0,0}$ , if this space is equipped with the norm

$$\|(\hat{v}, \hat{w})\|_N^2 = (\hat{v}, (-\hat{A} + \lambda_0)\hat{v})_{L^2} + (\hat{w}, \hat{w})_{L^2}.$$

Here  $\lambda_0 > 0$  is chosen large enough that  $-\hat{A} + \lambda_0$  is invertible. We have to verify that (A1) also holds in higher Sobolev spaces. As far as the space dependence is concerned, this is easily achieved by using norms involving the operator  $S$  (see [10-12]). However, the topology of  $Z$  also involves a  $\sigma$ -derivative. Note that

$$\frac{d}{d\sigma} A(\hat{v}, \hat{w}) = A\left(\frac{d\hat{v}}{d\sigma}, \frac{d\hat{w}}{d\sigma}\right) + \frac{dA}{d\sigma}(\hat{v}, \hat{w}).$$

Since  $\frac{dA}{d\sigma}$  is bounded from  $Z$  into  $W^{0,2} \times W^{0,1}$ , we can satisfy (A1) by choosing the same norm for the  $\sigma$ -derivative. The perturbation given by  $\frac{dA}{d\sigma}$  will not disturb the resolvent estimate required for the  $C_0$ -semigroup property.

#### REFERENCES

- [1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, *Comm. Pure Appl. Math.* 12 (1959), 623-727 and 17 (1964), 35-92.
- [2] B. Bernstein, E. A. Kearsley and L. J. Zapas, A study of stress relaxation with finite strain, *Trans. Soc. Rheology* 7 (1963), 391-410.
- [3] W. V. Chang, R. Bloch and V. W. Tschoegl, On the theory of viscoelastic behaviour of soft polymers in moderately large deformations, *Rheol. Acta* 15 (1976), 367-378.
- [4] C. M. Dafermos and J. A. Nohel, A nonlinear hyperbolic Volterra equation in viscoelasticity, *Amer. J. Math.*
- [5] M. Doi and S. F. Edwards, Dynamics of concentrated polymer systems, *J. Chem. Soc. Faraday* 74 (1978), 1789-1832 and 75 (1979), 38-54.
- [6] A. Douglis and L. Nirenberg, Interior estimates for elliptic systems of partial differential equations, *Comm. Pure Appl. Math.* 8 (1955), 503-538.
- [7] M. S. Green and A. V. Tobolsky, A new approach to the theory of relaxing polymeric media, *J. Chem. Phys.* 14 (1946), 80-100.
- [8] T. J. R. Hughes, T. Kato and J. E. Marsden, Well-posed quasi-linear second-order hyperbolic systems with applications to nonlinear elastodynamics and general relativity, *Arch. Rat. Mech. Anal.* 63 (1976), 273-284.
- [9] E. F. Infante and J. A. Walker, A stability investigation for an incompressible simple fluid with fading memory, *Arch. Rat. Mech. Anal.* 72 (1980), 203-218.
- [10] T. Kato, Linear evolution equations of "hyperbolic" type I, *J. Fac. Sci. Univ. Tokyo* 17 (1970), 241-258.
- [11] T. Kato, Linear evolution equations of "hyperbolic" type II, *J. Math. Soc. Japan* 25 (1973), 648-666.
- [12] T. Kato, Quasi-linear equations of evolution with application to partial differential equations, in: W. N. Everitt (ed.), *Spectral Theory of Differential Equations*, Springer Lecture Notes in Mathematics 448 (1975), 25-70.

- [13] A. Kaye, Co A Note 134, The College of Aeronautics, Cranfield, Bletchley, England 1962.
- [14] J. U. Kim, Global smooth solutions of the equations of motion of a nonlinear fluid with fading memory, Arch. Rat. Mech. Anal. 79 (1982), 97-130.
- [15] A. S. Lodge, A network theory of flow birefringence and stress in concentrated polymer solutions, Trans. Faraday Soc. 52 (1956), 120-130.
- [16] A. S. Lodge, Constitutive equations from molecular network theories for polymer solutions, Rheol. Acta 7 (1968), 379-392.
- [17] J. G. Oldroyd, On the formulation of rheological equations of state, Proc. Royal Soc. London A 200 (1950), 523-541.
- [18] M. Renardy, Singularly perturbed hyperbolic evolution problems with infinite delay and an application to polymer rheology, SIAM J. Math. Anal.
- [19] M. Renardy, Local existence theorems for the first and second initial-boundary value problem for a weakly non-Newtonian fluid, Arch. Rat. Mech. Anal.
- [20] M. Slemrod, A hereditary partial differential equation with applications in the theory of simple fluids, Arch. Rat. Mech. Anal. 62 (1976), 303-321.
- [21] C. A. Truesdell and W. Noll, The Non-Linear Field Theories of Mechanics, in: S. Flügge (ed.), Handbuch der Physik III/3, Springer 1965.
- [22] A. F. K. Ward and G. M. Jenkins, Normal thrust in dynamic torsion for rubberlike materials, Rheol. Acta 1 (1958), 110-114.
- [23] L. J. Zapas, Viscoelastic behaviour under large deformations, J. Res. Nat. Bur. Stand. 70A (1966), 525-532.

MR/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2530	2. GOVT ACCESSION NO. AD-A130503	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Local Existence and Uniqueness Theorem for a K-BKZ-Fluid		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Michael Renardy		8. CONTRACT OR GRANT NUMBER(s) MCS-7927062, Mod. 2. DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS  See Item 18 below		12. REPORT DATE June 1983
		13. NUMBER OF PAGES 15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709  National Science Foundation Washington, DC 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Quasilinear Hyperbolic Equations, Viscoelastic Liquids, Semigroup theory		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We consider the three-dimensional motion of a viscolastic liquid occupying all space. The constitutive law is assumed to be of the form suggested by Kaye [13] and Bernstein, Kearsley and Zapas [2]. An existence and uniqueness result for solutions of the initial value problem on sufficiently short time intervals is proved using Kato's theory of quasilinear hyperbolic equations.		

ND

ATE

LMED

-83

TIC